


RESEARCH

Open Access



Gestational hypoxia in late pregnancy differentially programs subcortical brain maturation in male and female rat offspring

E. Nicole Wilson³, Steve Mabry³, Jessica L. Bradshaw³, Jennifer J. Gardner^{1,3}, Nataliya Rybalchenko³, Rachel Engelland³, Oluwadarasimi Fadeyibi³, Oluwatobiloba Osikoya¹, Spencer C. Cushen^{1,2}, Styliani Goulopoulou^{1,4} and Rebecca L. Cunningham^{3*} 

Abstract

Background: Hypoxia is associated with pregnancy complications, such as preeclampsia, placental abruption, and gestational sleep apnea. Hypoxic insults during gestation can impact the brain maturation of cortical and subcortical pathways, such as the nigrostriatal pathway. However, the long-term effects of in utero hypoxic stress exposure on brain maturation in offspring are unclear, especially exposure during late gestation. The purpose of this study was to determine the impact of gestational hypoxia in late pregnancy on developmental programming of subcortical brain maturation by focusing on the nigrostriatal pathway.

Methods: Timed pregnant Long–Evans rats were exposed to chronic intermittent hypoxia or room air normoxia from gestational day (GD) 15–19 (term 22–23 days). Male and female offspring were assessed during two critical periods: puberty from postnatal day (PND) 40–45 or young adulthood (PND 60–65). Brain maturation was quantified by examining (1) the structural development of the nigrostriatal pathway via analysis of locomotor behaviors and the substantia nigra dopaminergic neuronal cell bodies and (2) the refinement of the nigrostriatal pathway by quantifying ultrasonic vocalizations (USVs).

Results: The major findings of this study are gestational hypoxia has age- and sex-dependent effects on subcortical brain maturation in offspring by adversely impacting the refinement of the nigrostriatal pathway in the absence of any effects on the structural development of the pathway. During puberty, female offspring were impacted more than male offspring, as evidenced by decreased USV call frequency, chirp USV call duration, and simple call frequency. In contrast, male offspring were impacted more than female offspring during young adulthood, as evidenced by increased latency to first USV, decreased simple USV call intensity, and increased harmonic USV call bandwidth. No effects of gestational hypoxia on the structural development of the nigrostriatal pathway were observed.

Conclusions: These novel findings demonstrate hypoxic insults during pregnancy mediate developmental programming of the cortical and subcortical pathways, in which male offspring exhibit long-term adverse effects compared to female offspring. Impairment of cortical and subcortical pathways maturation, such as the nigrostriatal pathway,

*Correspondence: rebecca.cunningham@unthsc.edu

³ Department of Pharmaceutical Sciences, UNT System College of Pharmacy, School of Pharmacy, University of North Texas Health Science Center, 3500 Camp Bowie Boulevard, Fort Worth, TX 76107, USA
Full list of author information is available at the end of the article



© The Author(s) 2022. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

may increase risk for neuropsychiatric disorders (e.g., mood disorders, cognitive dysfunction, brain connectivity dysfunction).

Highlights

- Brain maturation of the nigrostriatal pathway is sex- and age- dependent. Exposure to hypoxia in late pregnancy impacts brain maturation of the nigrostriatal pathway that can be observed during puberty and young adulthood.
- Gestational hypoxia impacted female offspring during puberty more than males, whereas it impacted male offspring during young adulthood more than females.
- These novel findings demonstrate that hypoxic insults during pregnancy mediate developmental programming of the cortical and subcortical pathways, in which male offspring exhibit long-term adverse effects compared to female offspring.
- Long-term adverse effects of gestational hypoxia in offspring can occur in the absence of pregnancy complications, especially if they occur within critical embryological developmental periods.

Keywords: Prenatal programming, Oxidative stress, Sex differences, Chronic intermittent hypoxia, Ultrasonic vocalizations, Substantia nigra

Background

Hypoxia or decreased oxygen during pregnancy is associated with many gestational complications, such as preeclampsia and gestational sleep apnea [1–5]. Gestational sleep apnea is a common late gestational (i.e., third trimester) hypoxic stressor that is observed in 26% of pregnant women [6, 7]. Gestational hypoxic stress can induce a multitude of long-term impairments across the physical, cognitive, and psychosocial domains, such as long-term memory problems [8–10] and motor impairments [5]. Perinatal insults at late gestation (i.e., third trimester) are specifically relevant to fetal brain development, because this is the period that cortical and subcortical maturation (e.g., neuroplasticity) occurs [11–16], such as the nigrostriatal pathway [11, 13, 17–19].

Male and female offspring respond differently to insults during gestation [20–22]. Prenatal insult paradigms in rodents showed that male offspring, and not female offspring, exhibited impairments of all maturation domains: psychosocial domain (elevated brain-regulated stress responses during adulthood [23, 24], social behavior [25]), cognitive domain (learning and memory deficits [21, 25]), and physical domain (decreased growth following weaning [23, 24] and impaired motor function [26]). Further, males are more sensitive to common insults that occur anytime during gestation, such as placental inflammation [27, 28], hypoxia [27, 28], term preeclampsia and eclampsia [26]. Although many studies have found that males may be more vulnerable to prenatal insults, recent studies demonstrated that prenatal insults can also negatively impact female offspring [26, 29, 30]. Animal studies that include females are not as extensively conducted,

and thus the impact of prenatal insults on female offspring is understudied [31–33].

Ultrasonic vocalizations (USVs) are dependent on multiple brain maturation domains, especially a functional nigrostriatal pathway in the brain [34–36]. USVs are calls emitted by rodents to other rodents to relay information [37–39]. They are innate behaviors and not learned [40, 41]. Fifty kilohertz (kHz) USVs were first described in the early 1970s, and they were associated with rodent mating behaviors [42]. However, 50 kHz USVs are not restricted to mating behaviors [43, 44], as they are emitted under several conditions, such as mother–pup retrieval, juvenile play interactions, same-sex interactions, and opposite-sex interactions [45–48]. Chirp and simple calls are the most common USV call type, especially during cage mate separation, and are generally associated with social coordination functions (i.e., re-establishing social contact) rather than reward or pleasure [41, 43, 49]. The more complex USVs, such as harmonic call types, are associated with sex behaviors, group housing, juvenile play, and other pleasurable behaviors (e.g., tickling) [50–54]. Larger USV bandwidths are associated with communicating to other rodents further away, indicating that the higher bandwidth may be used to compensate in a more expansive environment [55]. In addition to 50 kHz USVs, rodents vocalize at 22 kHz USVs under stressful or aversive conditions [56]. These 22 kHz USVs associated with distress or danger are generally long duration calls (> 300 ms) [56, 57].

The long-term effects of in utero hypoxic stress exposure on brain maturation in offspring are unclear, especially exposure during late gestation. Therefore, the purpose of this study was to determine the impact of late gestational hypoxia on the developmental programming of subcortical brain maturation by focusing on the nigrostriatal pathway. We used an intermittent hypoxia protocol between GD 15–19 to model gestational sleep apnea during the third trimester, which is observed in 26% of pregnant women [6, 7]. We focused on the nigrostriatal subcortical pathway that is established during GD 16–20 [11, 13, 17–19]. Brain maturation was quantified by examining (1) the structural development of the nigrostriatal pathway via analysis of locomotor behaviors and the dopaminergic neuronal cell bodies located in the substantia nigra and (2) the refinement of the nigrostriatal pathway by quantifying USVs that are sensitive to any nigrostriatal pathway impairments [34–36]. In addition to behavioral measures of the nigrostriatal pathway integrity, we examined oxidative stress, as the substantia nigra exhibits high basal levels of reactive oxygen species [58], by quantifying circulating oxidized proteins and oxidative stress-associated enzymatic activity in the substantia nigra. We hypothesized that exposure to hypoxia during late gestation would impair fetal brain maturation in a sex-dependent manner.

Methods

Animals

All experiments were conducted in agreement with the Guide for the Care and Use of Laboratory Animals of the National Institutes of Health and the ARRIVE guidelines. These protocols were approved by the Institutional Animal Care and Use Committee of the University of North Texas Health Science Center.

Dams

All experiments were conducted using timed pregnant Long-Evans rats (aged 8–10 weeks, Charles River, Wilmington, MA). Dams arrived at the University of North Texas Health Science Center animal facilities on gestational day 5–7 (term = 22–23 days). Animals were single-housed in a 12 h:12 h light/dark cycle with lights on at 0900 h and were provided food and water ad libitum. Two studies were conducted. The purpose of the first study ($n = 16$ dams) was to examine offspring behavior in response to late gestational hypoxia, while the purpose of the second study ($n = 12$ dams) was to characterize the effects of gestational hypoxia on fetal and placental biometrics. Animals were allowed to habituate for 1 week prior to the initiation of the chronic intermittent hypoxia (CIH) protocol.


Offspring

For study one, pup weights and measures of crown-rump length and abdominal girth were collected twelve to sixteen hours after birth. Sex was estimated by determination of anogenital distance, which was confirmed by the presence of gonads (testes or vagina) during the prepubertal stage. Litters were reduced to eight pups/litter and, when possible, to equal number of males and females. Pups were culled based on size, wherein heavier or lighter than the group average of pup offspring was removed. Weaning occurred on postnatal day (PND) 28. Immediately after weaning, dams were anesthetized with 2–3% isoflurane and euthanized via decapitation. Subsequently, four male and four female pups in each litter were randomly assigned to either pubertal ($n = 2$ /sex/litter) or young adult ($n = 2$ /sex/litter) groups for behavioral studies (Fig. 1). For study two, dams and fetuses were euthanized via decapitation on GD20 to collect fetal and placental biometrics [59].

Pubertal and young adult offspring (sexually naïve) were pair housed with their littermate of the same sex. Male offspring were moved to a separate room from the female offspring for the remainder of the study. Offspring were housed in rooms on a 12-h (h) reverse light cycle where lights were off at 0900 h. Reverse lighting allowed for behavioral testing to be conducted during offspring's wake phase of the circadian cycle. Food and water were provided to all animals ad libitum. Offspring body weights were obtained weekly after weaning. To acclimatize the offspring to operator handling and reduce stress responses during behavior testing, animals were handled daily, beginning approximately 10 days prior to the start of behavior testing. Pubertal and young adult offspring were anesthetized with 2–3% isoflurane and euthanized via decapitation on PND 48 or PND 66, respectively.

Gestational chronic intermittent hypoxia

To examine the impact of hypoxia at late gestation on offspring behaviors, timed-pregnant female rats were assigned to receive either chronic intermittent hypoxia (CIH) or normoxia (Study one: $n = 8$ /group; Study two: $n = 6$ /group) starting at 0900 h for 8 h during their sleep phase of the circadian cycle on gestational days 15–19. To induce CIH, the home cages of the pregnant rats were placed into Oxycycler chambers (76.2 × 50.8 × 50.8 cm, BioSpherix, Lacona, NY, USA). Animals were allowed to acclimatize to the chambers under normoxic (room air) conditions for a period of 4 days prior to starting the CIH protocol, in which oxygen was reduced from 21% (room air) to 10% and then returned to 21% using a 6-min cycle (10 cycles/h) over 8 h/day for a period of 5 days, as previously described [60–63]. Based on gestational exposure to hypoxia or normoxia, offspring from these dams were



included in one of the following groups (Fig. 1): male normoxic (MN), male CIH (MC), female normoxic (FN), and female CIH (FC).

Study two examined the effects of late gestation CIH on fetal and placental biometrics. This separate study used two different CIH protocols: (1) 6-min cycles (10 cycles/hour) over 8 h/day for a period of 5 days and (2) 8-min cycles (8 cycles/hour), as previously described [60, 63]. This resulted in three different groups: (1) CIH with 6-min cycles/hour ($n=3$), (2) CIH with 8-min cycles/hour ($n=3$), and normoxic controls that were simultaneously run with each of the CIH protocols ($n=6$). No statistical significance was found between the two CIH protocols on any pregnancy outcome measures, and thus these CIH groups were collapsed into one group ($n=6$).

Behavioral testing

Behavioral studies were conducted during puberty (PND 40–45) and young adulthood (PND 60–65) over two days during their wake period from 0945 to 1700 h. Behavior test orders were randomized. A minimum of 18 h

separated behavior testing in days one and two. Male animals were tested before females to avoid potential confounding effects of pheromones on behavior. Behavioral apparatuses were thoroughly cleaned with 70% ethanol between each animal. One hour following behavior testing on males, female behaviors were examined. All behavior studies were conducted under red lighting and recorded for later analysis.

Cage mate separation and ultrasonic vocalizations (USVs)


To conduct the cage mate separation test, cage mates were removed from their home cage and placed into a $50 \times 25 \times 30$ cm aquarium and allowed to explore for 2 min [41, 64]. After 2 min, one cage mate was removed and placed into a separate aquarium. Each aquarium was enclosed within a sound dampening chamber equipped with an Avisoft UltraSoundGate 116Hb (CM16/CPMA) condenser microphone (Avisoft Bioacoustics, Nordbahn, Germany). USVs emitted for five minutes in response to the cage mate separation were recorded using Avisoft-Recorder USGH software (Avisoft Bioacoustics, Nordbahn,

Germany). Separated cage mates will emit 50 kHz USVs, which are due to the separation and not the novelty of a new chamber [49]. This separation of cage mate protocol was used to avoid confounding effects of pheromones or hormones on USVs [44, 50–52, 65–67]. Recorded USVs were analyzed using Avisoft SASLab Pro (Version 5.2.12, Avisoft Bioacoustics, Nordbahn, Germany). USV spectrogram analyses were conducted using Avisoft recommended settings of fast Fourier transform (FFT)-length of 256 points and an overlap of 50%. Spectrograms had a frequency resolution of 0.977 kHz and a time resolution of 0.5 milliseconds (ms). A threshold between -40 and -50 decibels (dB) was chosen to remove background noise. If background noise interfered with USV classification, those USVs were not analyzed. Calls were manually selected within the spectrogram and categorized by call type [41, 45, 53, 64]. Experimenters analyzing USVs were blinded to the treatments.

USVs were characterized by total call frequency (number of calls over the 5-min interval), call type (chirp, simple, harmonic), latency to first call, duration, intensity (dB), and bandwidth (kHz). USV intensity and bandwidth are included in analysis as impairments of these properties are linked with impaired nigrostriatal pathway function [36]. Stress calls (22 kHz) were not observed in this study. Therefore, all USVs in this study were emitted at the 50 kHz frequency, which are USVs produced during conditions of reward, positive affect, or normal behaviors. USVs were separated into chirp, simple, and harmonic call types based on their duration and intensity (Fig. 2). Chirp calls were defined as low intensity flat USVs with a duration less than 0.02 s (20 ms), whereas simple calls were defined as flat USVs lasting longer than 0.02 s. Harmonic calls were defined as a single call with two distinct tones, creating a ‘harmony’ with varying duration and intensity. USVs calls that could not be classified as chirp, simple, or harmonic calls, were classified as other calls and included in the total USV call analysis (Additional file 1).

Open field test

Gross and fine motor activities were quantified using the San Diego Instruments Photobeam Activity System-Open field arena ($40.64 \times 40.64 \times 38.1$ cm) with a unidirectional ‘rearing’ bar located above a bi-directional main field bar, as previously described [62]. Open field behaviors of distance traveled in meters (m), number of unassisted rears, and number of total rears were tracked and analyzed for a period of 10 min. An unassisted rear was classified when the animal stood on its hindlimbs without touching a wall or requiring aid to stand fully erect.



Tissue sample collection and preparation

During the first 2 h of the animals’ wake phase, each animal was anesthetized with isoflurane (2–3%) and decapitated [60–63]. Trunk blood was collected in EDTA-coated tubes and centrifuged at $2000 \times g$ for 10 min at 4°C to collect plasma, which was stored at -80°C .

Each brain was quickly removed, flash frozen in phosphate-buffer saline (PBS), sliced into 1-mm coronal sections using a brain matrix, and brain nuclei containing the substantia nigra (-5.30 mm from Bregma) were isolated according to Paxinos and Watson’s brain atlas [68] using blunt 20-gauge needles attached to 1 mL syringes [60–62]. Micro-dissected brain samples were placed into microcentrifuge tubes to be stored at -80°C until protein analysis. Whole placental tissues were collected, weighed, and flash-frozen in liquid nitrogen before storing at -80°C on GD20. Prior to homogenization, the maternal decidua was removed from each placenta and the remaining tissue was used for immunoblotting.

Circulating oxidative stress

Circulating oxidative stress in plasma proteins was assayed using OxiSelect Advanced Oxidative Protein Products (AOPP) kit (Cell Biolabs, Inc., San Diego, CA) according to our previously published protocol [60]. This assay measures the micromolar (μM) concentration of all oxidized proteins in the plasma relative to a known standard. Chloramine in the kit reacts with oxidized proteins to produce a color change, which can be read at 340 nm (nm).

Western blot

Frozen tissue samples were homogenized in RIPA lysis buffer (VWR, cat # N653) containing (per 0.5 ml): 2.5 μl HaltTM protease and phosphatase inhibitor (Thermo Scientific, cat # 78442), 1 μl 0.5 M ethylenediaminetetraacetic acid (EDTA, Sigma-Aldrich), and 1 μl 0.5 mM dithiothreitol (DTT, Sigma-Aldrich), as previously described [60]. For total protein concentration determination, protein levels in the homogenate were determined using Pierce BCA Protein Assay (Thermo Fisher, cat # 23225). Samples were denatured with β -mercaptoethanol and boiled at 100 °C for 5 min. Equal volumes containing 15–30 μg protein were loaded into a Bio-Rad 4–15% polyacrylamide gel. Electrophoresis was conducted at 25 milliamps (mA) in a Tris–glycine running buffer followed by overnight transfer at 4 °C onto a PVDF membrane at 50 mA. Following 30 min washing, membranes were blocked for 30 min with 5% non-fat milk in Tris-based saline (TBS)–Tween (TBS-T) at room temperature. Membranes were then transferred to 1% nonfat milk TBS-T solutions containing specific primary antibodies (anti-tyrosine hydroxylase, Millipore AB318 1:1000; Anti-Spectrin, EMD Millipore MAB1622 1:1000) and incubated overnight at 4 °C. Primary antibody against beta actin (GeneTex GTX 629630, 1:5000) was incubated for 1 h at room temperature. Afterwards, membranes were washed in 10 min increments for 30 min, and then incubated in 1% milk TBS-T with secondary antibody solutions (HRP-conjugated goat anti-mouse, Invitrogen 31430, 1:5000) at room temperature for 1 h. Protein bands were visualized using West Pico enhanced chemiluminescence detection assay (Thermo Scientific, cat # 34580) on a Syngene G:Box system using FlourChem HD2 AIC software as previously described [69]. NIH Image J software (version 1.48v) was used to quantify band densitometry, and values were normalized to beta actin.

Statistical analyses

Significance was defined as $p < 0.05$. Statistical analyses were conducted in Prism Version 8.4 (GraphPad, San Diego, CA). Outliers were identified and removed using

the ROUT method (coefficient $Q = 1\%$). Data distribution was tested using the Shapiro–Wilk test. For analysis of pubertal (PND 48) and young adult (PND 66) offspring outcomes, data with non-Gaussian distributions were normalized using square root transformation. Litter size, fetal and placental weights, and placental protein content were compared using an unpaired t test, while gestational length was compared using a Mann–Whitney U test. A two-way ANOVA (condition, sex) followed by Sidak's or Tukey's posthoc test was used to analyze neonatal biometrics and pubertal and young adult behaviors and protein expression. Data are presented as mean with S.E.M, unless otherwise indicated.

Results

Gestational hypoxia during late gestation did not affect maternal, fetal, or offspring body weights

All timed pregnant rats were weighed (203–276 g) upon arrival at the University of North Texas Health Science Center. No differences were observed in dam weights prior to initiation of CIH (normoxia: 250.8 ± 5.7 g, CIH: 255.9 ± 6.47 g, $p > 0.05$). In study one, dams were sacrificed during the postpartum period, 28 days following CIH. In study two, dams were sacrificed on GD20, approximately 16 h following the last bout of CIH. Maternal postpartum body weights at euthanasia did not differ between groups (normoxia: 284.14 ± 16.46 g, CIH: 282.45 ± 19.04 g; $p > 0.05$). Similarly, CIH exposure did not impact maternal body weights on GD20 (normoxia: 344.67 ± 5.30 g, CIH: 328.81 ± 9.87 g; $p > 0.05$).

Gestational CIH did not affect litter size (normoxia: 11.5 ± 1.8 , CIH: 9.7 ± 1.6 ; $p > 0.05$), number of resorptions (normoxia: 0.0 ± 0.0 vs. CIH: 0.5 ± 1.2 ; $p > 0.05$), fetal weights (normoxia: 2.48 ± 0.07 g, CIH: 2.37 ± 0.05 g; $p > 0.05$), placental weights (normoxia: 0.49 ± 0.02 g, CIH: 0.53 ± 0.02 g; $p > 0.05$), or placental efficiency as measured by fetoplacental weight ratios (normoxia: 5.07 ± 0.47 vs. CIH: 4.57 ± 0.54 ; $p > 0.05$) at GD20. Similarly, there were no differences in body weights between neonates from

Table 1 Neonatal biometrics after birth

		Treatment	
		Normoxia	CIH
Neonatal weight (g)	Male	6.55 ± 0.23	6.63 ± 0.21
	Female	6.43 ± 0.20	6.42 ± 0.23
Crown-rump length (cm)	Male	4.08 ± 0.08	4.05 ± 0.05
	Female	4.06 ± 0.08	4.06 ± 0.08
Abdominal girth (cm)	Male	4.82 ± 0.12	4.75 ± 0.12
	Female	4.70 ± 0.10	4.74 ± 0.06

Means \pm SEM. Two-way ANOVA with Sidak's multiple comparison test ($n = 8$ /group). All comparisons had p -values > 0.05 . CIH chronic intermittent hypoxia

CIH-exposed pregnancies compared to normoxic pregnancies (Table 1). Gestational CIH did not affect pubertal (PND48) offspring body weights (Males, normoxia: 295.9 ± 6.72 g, CIH: 285.6 ± 7.70 g; $p > 0.05$, Females, normoxia: 220.4 ± 3.46 g, CIH: 222.5 ± 5.1 g; $p > 0.05$) or young adult (PND66) offspring body weights (Males, normoxia: 438.3 ± 5.98 g, CIH: 460.6 ± 8.35 g; $p = 0.098$; Females, normoxia: 287.5 ± 6.24 g, CIH: 276.8 ± 7.00 g; $p > 0.05$).

Gestational length was greater in the CIH group compared to control, but this difference was not statistically significant (median (IQR), normoxia: 22 (22–23) days; CIH: 23 days (23–23); $p = 0.06$). There were no differences in number of live births between groups (normoxia: 12.38 ± 1.27 , CIH: 10.63 ± 1.05 ; $p > 0.05$).

Gestational hypoxia reduced total calls emitted by pubertal offspring and increased latency to first call in young adult male offspring

Gestational hypoxia reduced the frequency of total USVs (all call types) emitted by pubertal offspring ($p = 0.002$, Fig. 3A, Additional file 1). Female pubertal offspring from normoxic pregnancies emitted significantly more calls compared to male pubertal offspring exposed to gestational CIH. There was no effect of gestational CIH on total number of calls emitted by male or female young adult offspring ($p > 0.05$, Fig. 3B). However, young adult offspring from normoxic pregnancies emitted fewer calls compared to pubertal offspring from normoxic pregnancies ($p = 0.004$), with female young adult offspring emitting significantly fewer calls compared to female pubertal offspring ($p = 0.018$). Although gestational CIH reduced total call frequency in pubertal offspring, there was no effect of gestational CIH on latency to first call in male




Fig. 3 Total call frequency and latency to first call. Total call frequency (A, B) and latency to first call (C, D) produced by pubertal (A, C) and young adult (B, D) male and female offspring exposed to normoxia or chronic intermittent hypoxia (CIH) in utero. Analyzed by Two-way ANOVA with Tukey's multiple comparisons test, $n = 9-17$ /group. Frequency = number of calls per 5-min interval; s = seconds. * = main effect of CIH; & = Tukey's comparison. *, & = $p < 0.05$

or female pubertal offspring ($p=0.089$, Fig. 3C). In young adults, the effect of gestational CIH on latency to first call was dependent on sex ($p=0.018$), with an increase in latency to first call in CIH-exposed young adult male offspring compared to CIH-exposed young adult females (Fig. 3D).

Gestational hypoxia impacted chirp calls in pubertal and young adult offspring

Since gestational CIH decreased the total number of offspring vocalizations, we categorized the calls to determine if these differences were attributed to a specific call type. Chirp calls were defined as low-intensity USVs with a duration less than 20 ms (Fig. 2). Gestational hypoxia reduced frequency of chirp calls emitted by pubertal offspring ($p=0.010$, Fig. 4A, Additional file 1), and this effect was also present in young adult offspring ($p=0.010$, Fig. 4B). The effects of gestational CIH on chirp call frequency in pubertal offspring were dependent on sex ($p=0.021$, Fig. 4A). Specifically, female pubertal offspring exposed to gestational CIH emitted fewer chirp calls compared to pubertal females from normoxic pregnancies. There was no effect of gestational CIH on frequency of chirp calls emitted by pubertal males compared to age-matched males from normoxic pregnancies. Moreover, reductions in chirp calls were not observed in young adult females exposed to gestational CIH compared to young adult females from normoxic pregnancies (Fig. 4B). Gestational CIH did not affect chirp call duration in young adult offspring ($p>0.05$, Fig. 4D) but reduced the duration of chirp calls emitted by pubertal offspring ($p=0.032$, Fig. 4C). Although gestational CIH affected chirp call frequency and duration, CIH exposure did not impact the intensity of chirp calls emitted by pubertal offspring ($p>0.05$, Fig. 4E). Moreover, gestational CIH did not affect chirp call intensity in young adult offspring ($p>0.05$, Fig. 4F). In addition, gestational CIH did not affect chirp call bandwidth in pubertal offspring ($p>0.05$, Fig. 4G). Nonetheless, there was a sex effect on chirp call bandwidth in young adult offspring, with males emitting chirp calls of significantly increased bandwidth compared to females ($p=0.005$, Fig. 4H). Specifically, young adult males exposed to gestational CIH emitted chirp calls of greater bandwidth than CIH-exposed young adult females.

Pubertal female offspring exposed to gestational hypoxia emit fewer simple calls

Simple calls were defined as USVs of greater than 20 ms duration that can be visibly seen as a straight line (Fig. 2). Gestational CIH reduced simple calls emitted by pubertal

offspring ($p=0.028$), and this effect was dependent on sex ($p=0.022$, Fig. 5A, Additional file 1). Precisely, CIH-exposed pubertal females emitted fewer simple calls than female pubertal offspring from normoxic pregnancies. Young adult females emitted more simple calls than young adult males ($p=0.024$); however, there was no effect of gestational CIH on the production of simple calls by young adult offspring ($p=0.914$, Fig. 5B). There were no effects of gestational CIH on simple call duration in pubertal ($p>0.05$, Fig. 5C) or young adult offspring ($p>0.05$, Fig. 5D). Although not statistically significant, female young adult offspring emitted simple calls with greater duration than male young adult offspring ($p=0.063$, Fig. 5D).

There was no effect of gestational CIH on intensity of simple calls emitted by pubertal offspring ($p>0.05$, Fig. 5E). However, CIH exposure reduced the intensity of simple calls emitted by young adult offspring ($p=0.022$), and this effect was dependent on sex ($p=0.005$, Fig. 5F). Specifically, CIH-exposed young adult males emitted simple calls with lesser intensity (decreased decibels) compared to young adult offspring from normoxic pregnancies. Similar to pubertal simple call duration and intensity, gestational CIH had no effect on the bandwidth of simple calls emitted by pubertal offspring ($p>0.05$, Fig. 5G). Moreover, there was no effect of gestational CIH on bandwidth of simple calls emitted by young adult offspring ($p>0.05$, Fig. 5H). Yet, young adult males emitted simple calls with greater bandwidths compared to young adult female offspring ($p=0.034$).

Gestational hypoxia increased complexity of harmonic calls in young adult males

Harmonic calls are USVs with two separate, yet identifiable tones, creating a 'harmony' (Fig. 2). In pubertal offspring, there was no effect of gestational CIH on frequency of harmonic calls ($p>0.05$), harmonic call duration ($p>0.05$), harmonic call intensity ($p>0.05$), or harmonic call bandwidth ($p>0.05$, Fig. 6A, Additional file 1). Similarly, gestational CIH did not affect young adult offspring frequency of harmonic calls ($p>0.05$), harmonic call duration ($p>0.05$), or harmonic call intensity ($p>0.05$). However, young adult males produced harmonic calls with greater bandwidths compared to young adult female offspring ($p=0.007$, Fig. 6B). Additionally, harmonic call bandwidth was greater in CIH-exposed young adult offspring compared to young adult offspring from normoxic pregnancies ($p=0.004$). Specifically, CIH-exposed male young adults emitted harmonic calls with greater bandwidths than young adult females from normoxic pregnancies.




Fig. 4 Characterization of pubertal and young adult offspring chirp calls. Frequency (A, B), duration (C, D), intensity (E, F), and bandwidth (G, H) of chirp calls emitted by pubertal (A, C, E, G) and young adult (B, D, F, H) male and female offspring exposed to normoxia or chronic intermittent hypoxia (CIH) in utero. Analyzed by Two-way ANOVA with Tukey's multiple comparisons test, $n = 8-14$ /group. Frequency = number of calls per 5-min interval, ms = milliseconds, dB = decibels, kHz = kilohertz. * = main effect of CIH; # = main effect of sex; & = Tukey's comparison; *, #, & = $p < 0.05$




Fig. 5 Characterization of pubertal and young adult offspring simple calls. Frequency (A, B), duration (C, D), intensity (E, F), and bandwidth (G, H) of simple calls emitted by pubertal (A, C, E, G) and young adult (B, D, F, H) male and female offspring exposed to normoxia or chronic intermittent hypoxia (CIH) in utero. Analyzed by Two-way ANOVA with Tukey's multiple comparisons test, $n = 4-10$ /group. Frequency = number of calls per 5-min interval, ms = milliseconds, dB = decibels, kHz = kilohertz. * = main effect of CIH; # = main effect of sex; & = Tukey's comparison; *, #, & = $p < 0.05$




Fig. 6 Bandwidth of harmonic calls emitted by pubertal and young adult offspring. Bandwidth of harmonic calls produced by (A) pubertal and (B) young adult male and female offspring exposed to normoxia or chronic intermittent hypoxia (CIH) in utero. Analyzed by Two-way ANOVA with Tukey's multiple comparisons test, $n = 5-12/\text{group}$. kHz = kilohertz. * = main effect of CIH; # = main effect of sex, & = Tukey's comparison; *, #, & = $p < 0.05$

Gestational hypoxia did not impact offspring motor function

Gross and fine motor skills were assessed using an open field behavior test. Gross motor function was quantified

by examining distanced traveled (meters) and total rearing behaviors, while fine motor function was assessed by observing unassisted rearing behaviors during a 10-min test. No significant differences were observed in pubertal




Fig. 7 Distance traveled and total rearing behavior by pubertal and young adult offspring during modified open field behavior test. Total distance traveled (A, B) and total rears (C, D) by pubertal and young adult male and female offspring exposed to normoxia (A, C) or chronic intermittent hypoxia (CIH, B, D) in utero. Analyzed by Two-way ANOVA with Tukey's multiple comparisons test, $n = 9-16/\text{group}$. kHz = kilohertz. * = main effect of CIH, & = Tukey's comparison; *, #, & = $p < 0.05$

animals in distance traveled regardless of sex or exposure to gestational CIH ($p > 0.05$). Similarly, no significant differences were observed in distance traveled by young adult males or females ($p > 0.05$). Distance traveled was significantly greater in young adult offspring from normoxic pregnancies compared to pubertal offspring from normoxic pregnancies ($p < 0.0001$), and this effect was significant for males and females (Fig. 7A). Distance traveled was significantly greater in young adult offspring from CIH exposed pregnancies compared to pubertal offspring from CIH exposed pregnancies ($p = 0.0006$). This effect was significant for males and not for females (Fig. 7B).

Total rearing behaviors were also assessed to examine gross motor behavior. Total rears were significantly greater in young adult offspring from normoxic pregnancies than pubertal offspring from normoxic pregnancies ($p = 0.031$) (Fig. 7C). Gestational CIH did not impact total rearing behavior, regardless of sex or age ($p > 0.05$) (Fig. 7D). When assessing fine motor function, no impact of age or sex was observed on unassisted rears in offspring from normoxic pregnancies (MN-puberty: 12.42 ± 3.32 , MN-young adult: 14.13 ± 2.09 , FN-puberty: 7.10 ± 2.12 , FN-young adult: 13.30 ± 2.01 ; $p > 0.05$). Similarly, there was no effect of gestational CIH on unassisted rears by young adults or pubertal rats (MC-puberty: 9.13 ± 1.95 , MC-young adult: 12.91 ± 1.91 , FC-puberty: 9.00 ± 1.84 , FC-young adult: 7.44 ± 2.44 ; $p > 0.05$).

Gestational hypoxia increased oxidative stress in young adult offspring

Since we observed CIH-induced deficits in pubertal and young adult offspring USVs, we examined markers of oxidative stress within the circulation and in specific brain regions associated with USVs and motor function. Circulating oxidative stress (AOPP) was not increased in pubertal offspring regardless of sex or exposure to gestational CIH ($p > 0.05$, Fig. 8A). Circulating oxidative stress was not increased in young adult offspring exposed to gestational CIH compared to age matched offspring from normoxic pregnancies, ($p = 0.067$, Fig. 8B).

In many neurological pathologies, damage to neurons precedes onset of symptoms. Abnormal proteolysis of cytoskeletal proteins, such as spectrin, by proteases including calpains and caspases contributes to neuronal cell damage or even death [70, 71]. Thus, we examined calpain cleavage (oxidative stress-associated mechanisms) and caspase-3 cleavage (apoptosis-associated mechanisms) of spectrin in the substantia nigra, a brain region associated with rodent USVs and motor function (Table 2) [34–36]. Within the substantia nigra of pubertal and young adult offspring, no significant differences were observed in calpain cleavage of spectrin regardless of sex or exposure to gestational CIH ($p > 0.05$). Likewise, there were no observed effects ($p > 0.05$) of sex or gestational CIH on caspase-3 cleavage of spectrin, an indicator of apoptosis [70, 71], in the substantia nigra of pubertal and young adult offspring. Since prior studies found that loss of dopaminergic neurons in the substantia nigra impaired




Table 2 Protein expression analysis of damage-associated markers in substantia nigra and placenta

		Substantia nigra				Placenta	
		Puberty		Young adult		GD20	
		Normoxia	CIH	Normoxia	CIH	Normoxia	CIH
Calpain	M	163.3 ± 28.44	180.6 ± 18.09	204.3 ± 48.88	148.4 ± 38.99	123.2 ± 21.43	115.9 ± 22.57
	F	137.4 ± 31.07	165.2 ± 18.14	168.6 ± 17.66	232.2 ± 45.08		
Caspase-3	M	51.68 ± 12.72	60.75 ± 13.19	75.23 ± 24.04	41.47 ± 11.15	46.68 ± 8.11	47.96 ± 13.00
	F	46.55 ± 18.38	40.16 ± 9.60	70.15 ± 12.70	66.31 ± 14.90		
TH	M	108.9 ± 25.36	143.0 ± 53.12	106.9 ± 40.22	154.4 ± 47.87	Not examined	
	F	155.4 ± 43.11	98.58 ± 32.69	69.44 ± 22.75	151.9 ± 39.41		

Values represent percentage of Beta Actin expression and are displayed as means ± SEM. Two-way ANOVA with Tukey's multiple comparison test for substantia nigra comparisons ($n = 5-6/\text{group}$). Unpaired t test for placenta comparisons ($n = 6/\text{group}$). No significant differences in any comparisons ($p > 0.05$). *GD* gestational day, *CIH* chronic intermittent hypoxia, *M* male, *F* female, *TH* tyrosine hydroxylase

USVs [36, 72, 73], we assessed if gestational CIH at late gestation impacted dopaminergic neuron expression in the substantia nigra. We used a marker of dopaminergic neurons, tyrosine hydroxylase, to ascertain whether dopaminergic neurons within the substantia nigra were damaged (Table 2). There were no significant differences in tyrosine hydroxylase expression within the substantia nigra of pubertal and young adult offspring, regardless of sex or gestational CIH ($p > 0.05$). To assess whether gestational CIH during late gestation induced damage to placental tissue, we also examined spectrin cleavage by calpain and caspase-3 within placentas isolated on GD20 (Table 2). There was no effect of CIH exposure on calpain or caspase-3 cleavage of placental spectrin ($p > 0.05$).

Discussion

The hypothesis for this study was that exposure to hypoxia during late gestation would impair fetal brain maturation in a sex-dependent manner. The major findings are (1) brain maturation of the nigrostriatal pathway is sex- and age-dependent in rats and (2) hypoxia during late gestation impacts rat brain maturation of the nigrostriatal pathway that can be observed during puberty and young adulthood in offspring.

We observed that hypoxic stress during late pregnancy impacted brain maturation of the nigrostriatal pathway that can be observed during puberty and young adulthood but did not impact the structural development of the nigrostriatal pathway. Specifically, gestational CIH impacted brain maturation in female offspring more than male offspring during puberty: (a) decreased chirp USV call frequency and (b) decreased simple USV call frequency. In contrast, brain maturation in male offspring during young adulthood was more impacted than female offspring: (a) increased latency (or time) to first USV and (b) decreased simple call intensity. Interestingly, late gestational CIH quantitatively impaired (ability to make and

sustain a call) the emission of USVs by females only during puberty, whereas CIH qualitatively impaired (USV call characteristics) the emission of USVs by adult males. It should be noted that the meaning of the different USVs is still relatively unknown. In this study, we used the cage mate separation protocol to elicit USVs, which is associated with re-establishing social contact. It is possible that late gestational hypoxia could have different effects on USV call characteristics in different environments, such as during sex behavior interactions. Although this is the first study to examine the impact of gestational CIH (GD 15–19) on USVs emitted by offspring during puberty and young adulthood, our findings are consistent with studies in humans that found prenatal hypoxic stress has more long-term effects in male offspring than female offspring [27, 28]. However, the literature on sex differences in response to gestational hypoxia in rodents is unclear, wherein there is a range of responses from differences in pregnancy outcomes, neonatal biometrics, and offspring physiology (e.g., oxidative stress, motor function).

Most gestational hypoxia studies use a long-term hypoxia protocol (> 12 days exposure) that encompasses all three gestational stages of pregnancy (Table 3), which is consistent with brain development in humans [74–76]. Brain developmental stages include (1) stage one wherein the neural progenitor cells become differentiated and the neural tube is formed [76, 77], (2) stage two when the cortical and subcortical regions of the brain are established with neural production [11–16, 77], and (3) stage 3 when maturation of brain cortical and subcortical pathways occur [16, 18, 77]. The effects of long-term gestational hypoxia on pregnancy outcomes or neonatal biometrics are generally dependent on the hypoxia protocol used (sustained hypoxia versus intermittent hypoxia cycles), in which sustained hypoxia is more likely to impact pregnancy outcomes and neonatal biometrics (Table 3). Similarly, the sustained hypoxia protocol negatively affects

Table 3 Comparison of gestational hypoxia parameters on offspring outcomes

Long-term gestational hypoxia (≥ 12 days): brain developmental stages 1–3						
Species	Gestational exposure	O ₂ (%)	Hypoxia (hours/day)	Hypoxic cycles from 21% O ₂	Impact on offspring	Citations
Rat (SD)	GD 5–21; 17 days	10%	8 h sleep phase	20 cycles/hour	<i>Age effects:</i> At birth: Decreased birth weight Puberty: No effect on body weight Adult: No effect on body weight	[115]
Rat (SD)	GD 3–19; 17 days	4%	8 h sleep phase	60 cycles/hour	<i>Age effects:</i> At birth: No effect on litter size or sex distribution Puberty: Decreased body weight Adult: Decreased body weight, increased myocardial SOD1 <i>Sex differences:</i> Adult: Increased myocardial lipid peroxidation in hypoxic males	[116]
Rat (SD)	GD 10–21; 12 days	10.5%	8 h sleep phase	15 cycles/hour	<i>Age effects:</i> At birth: No effect on litter size or body weight. No oxidative stress in placenta or fetal brains. Increased maternal separation-induced USVs at PND 4 Adult: No effects on open field locomotor behavior <i>Sex differences:</i> At birth: Increased maternal separation-induced harmonic calls in hypoxic females	[117]
Rat (SD)	GD 7–21; 15 days	10%	3 h	Sustained	<i>Age effects:</i> At birth: Decreased birth weight Puberty: No effect on body weight Adult: No effect on body weight	[118]
Rat (SD)	GD 7–21; 15 days	10%	3 h	Sustained	<i>Age effects:</i> At birth: No effect on litter size, decreased birth weight	[119]
Rat (Wistar)	GD 6–20; 15 days	13%	24 h	Sustained	<i>Age effects:</i> At birth: Increased placental weight, no effects on birth weight, fetal biometrics, litter size, or sex ratio. No oxidative stress or lipid peroxidation in hypoxic placentas	[120]
Mouse (C57BL/6 J)	GD 6–18; 13 days	14%	24 h	Sustained	<i>Sex differences:</i> Adult: Increased oxidative stress in hypoxic males but decreased oxidative stress in hypoxic females	[121]
Mid-term gestational hypoxia (6–9 days): brain developmental stages 2–3						
Species	Gestational exposure	O ₂ (%)	Hypoxia (hours/day)	Hypoxic cycles from 21% O ₂	Impact on offspring	Citations
Rat (SD)	GD 15–20; 6 days	11%	24 h	Sustained	<i>Sex differences:</i> Adult: Decreased body weight in hypoxic males	[122]
Mouse (C57BL/6 J)	GD 10–18; 9 days	10.5%	24 h	Sustained	<i>Age effects:</i> At birth: Decreased litter size, crown to rump length, and abdominal circumference. No effects on placental weight or placental oxidative stress	[123]

Table 3 (continued)**Short-term gestational hypoxia (2–5 days):
brain developmental stage 3**

Species	Gestational exposure	O ₂ (%)	Hypoxia (hours/day)	Hypoxic cycles from 21% O ₂	Impact on offspring	Citations
Rat (Long-Evans)	GD 15–19; 5 days	10%	8 h—sleep phase	10 cycles/hour	<i>Age effects:</i> At birth: No effect on litter size, body weights, placental weights, crown–rump length, or abdominal girth. Increased gestational length Puberty: Decreased USV calls. No effect on locomotor behavior (distance traveled or rearing) Adult: Increased bandwidth of harmonic USVs and increased oxidative stress. No effect on locomotor behaviors (distance traveled or rearing) <i>Sex differences:</i> Puberty: Decreased chirp USVs, chirp duration, decreased simple USVs in hypoxic females Adult: In males—increased latency to first USV and harmonic USV bandwidth with decreased chirp USVs and simple call USV intensity. In females—increased simple USV duration	Current Study
Mouse (CD-1)	GD 14–17; 4 days	11%	24 h	Sustained	<i>Age effects:</i> At birth: Decreased birth weight. No change in placental weights or litter size	[124]

**Acute gestational hypoxia (< 1 day):
brain developmental stage 3**

Species	Gestational exposure	O ₂ (%)	Hypoxia (hours/day)	Hypoxic cycles from 21% O ₂	Impact on offspring	Citations
Mouse (C57BL/6 J)	GD 17	9%	2 h	Sustained	<i>Age effects:</i> At birth: Increased brain weight Puberty: Increased volume of ventral tegmental area, pontine central grey matter, and cerebellar white matter (PND 23) Adult: No differences in brain volume	[125]
Mouse (C57BL/6 J)	GD 17.5	5%	2, 4, 6, or 8 h	Sustained	<i>Age effects:</i> At birth: No difference in litter size or gestational length. Decreased body weight that normalized at PND 8. No differences in brain apoptosis (cortex, basal ganglia, white matter) Adult: No differences in body weight, lateral ventricle size, anterior cingulate cortical thickness, or corpus callosum white matter. Decreased hindlimb strength	[126]

SD Sprague–Dawley, *GD* gestational day, *PND* postnatal day, *USV* ultrasonic vocalizations, *SOD1* superoxide dismutase-1

pregnancy outcomes and neonatal biometrics in shorter hypoxia protocols (6–9 days) that are restricted to brain development stages of 2–3. Based on findings that show no effects of acute sustained hypoxia protocols (< 1 day) during the final day of gestation on pregnancy outcomes and neonatal biometrics, development stages 2–3 are hypothesized to be the most important developmental stages for pregnancy outcomes and neonatal biometrics. Using an intermittent hypoxia cycling protocol that was restricted to GD 15–19 during brain developmental stage 3, our study focused on brain pathway maturation

of the nigrostriatal subcortical pathway in offspring in the absence of negative effects on the process of placentation and having minimal impact on pregnancy outcomes and neonatal biometrics (Table 3).

Brain regions that are sensitive to prenatal hypoxia include the cortex, basal ganglia (includes the nigrostriatal pathway), and white matter in children [78]. The nigrostriatal pathway is integral for many behaviors, such as motor function, cognition, social interactions, mood (anxiety/depression), reward, and attention [79–81]. Therefore, impairments of the nigrostriatal pathway can

have long-term consequences, such as many neuropsychiatric disorders (e.g., Parkinson's disease [78, 82], addiction [78], cognitive dysfunction [78], autism spectrum disorders [78, 79], mood disorders [78, 79, 81], and attention deficit hyperactivity disorders [80]). These relationships have also been observed in clinical studies examining children exposed to gestational hypoxia (pre-eclampsia, acute perinatal hypoxic injury, and gestational sleep apnea [1–5]). Preeclampsia is associated with autism spectrum disorders, attention deficit disorders, schizophrenia, mood disorders, developmental delay, cognitive dysfunction, stroke, cerebral palsy, and brain connectivity [83–94]. Similarly, gestational sleep apnea is associated with neuropsychiatric outcomes in children, such as social interactions [95]. Interestingly, gestational sleep apnea impacts male children more than female children with respect to social interactions, cognitive function, and communication [96], which is consistent with the findings from our study. Moreover, the hypoxic protocol used in the current study is consistent with gestational sleep apnea, in which the hypoxic stress is intermittent and only present during the sleep period.

In addition to gestational CIH effects, we observed age- and sex-related developmental effects on brain maturation. Young adult males from normoxic pregnancies emitted USV calls (all call types) at greater bandwidths than young adult females, which is in agreement and extends prior findings showing male mice emitted larger bandwidths in complex calls than female mice [67]. Further, we showed that pubertal rats emitted more USVs than young adult rats. Prior studies have not compared pubertal USV call frequencies to young adult USV call frequencies, much less USV call frequencies in females. However, a recent study comparing USV call frequency between adult male C57BL/6 J mice at various ages (PND 56, PND 140, and PND 210) found that USV call frequencies declined with age in sexually naïve male mice, whereas this relationship with age was not observed in sexually experienced male mice [97]. Although pubertal C57BL/6 J mice (~PND 30) were not included in this study [97], it is probable that age is the main factor for the observed decline in USV call frequency observed in our findings between sexually naïve young adult rats and pubertal rats.

Interestingly, our study demonstrates young adult females vocalize more simple USV calls compared to males, with no sex differences observed in other USV call frequencies (total USVs, chirp USVs, harmonic USVs). This is in contrast to findings from several other studies that found males emit more USVs than females [45, 65, 67, 98, 99]. It should be noted that these other studies were conducted under different parameters than the current study, in which USVs were quantified during

male–female interactions. Changing the parameters or the environment that USVs are collected can dramatically impact USVs. Indeed, no sex differences in USVs are observed under conditions of same-sex interactions in mice [100]. Female mice emit more USVs in the presence of novel female conspecifics than familiar female conspecifics [101]. To remove the potential confounds of sexual experience, hormonal conditions, or novelty from conspecific animals that can independently impact USVs [44, 50–52, 65–67], we used the cage mate separation protocol to elicit USVs [49]. The cage mate separation protocol to elicit USVs has only been used in male rats [49, 54], but our results show that this protocol is sufficient to also elicit USVs in female rats. This cage mate separation protocol resulted in unmasking a previously unknown sex difference in which young adult female rats vocalize more simple USVs than young adult male rats.

Additionally, we demonstrated sex- and age-dependent differences in motor activity. Specifically, young adult male and female rats exhibited greater locomotor activity (distance traveled) than pubertal male and female rats. Young adult females displayed increased total rearing behaviors compared to pubertal females. Our results on motor function are consistent with prior reports in the literature. Several studies have reported increased total rearing behaviors in young adult female rats compared to pubertal female rats, young adult male rats, and pubertal male rats [102–105].

In contrast to the literature on total rearing behaviors, previous studies on sex differences in locomotion are equivocal. Several studies show increased locomotor activity in young adult male and female rats compared to pubertal male and female rats [102, 106–109], which is consistent with our results. Other studies have shown increased locomotor activity in female rats compared to male rats, whereas others reported no sex differences in locomotor activity in rats that range in age from PND 30–PND 80 [110, 111]. These differences in locomotor activity could be due to environmental factors, such as novelty of the open field arena and the size of the open field arena [108, 112, 113]. Previous studies found increased locomotor activity in young adult (PND 60) male rats compared to pubertal (PND 42) male rats when animals were placed into a novel open field arena, but this effect was not observed in rats that were habituated to the open field arena prior to testing to decrease spontaneous exploratory behaviors [108]. Furthermore, the size of the open field arena has been shown to impact the ability to observe sex differences in locomotor behavior. Increased locomotion by female rats compared to male rats has been observed in large open field arenas (129 × 120 × 60 cm) [112]. However, other studies have shown no sex differences in locomotor activity

when using smaller open field arenas (70 × 70 × 70 cm or 54.5 × 80 × 33 cm) [113, 114]. It should be noted that in our study the rats were placed into a novel open field arena (40.64 × 40.64 × 38.1 cm), which may be why we observed age-associated changes in distance traveled locomotor activity but not sex-associated changes.

Perspectives and significance

The major findings of this study are that gestational hypoxia has age- and sex-dependent effects on subcortical brain maturation in offspring, such as the nigrostriatal pathway. During puberty, female offspring were impacted more than male offspring. In contrast, male offspring were impacted more than female offspring during young adulthood, indicating long-term adverse effects in male offspring compared to female offspring. Impairment of cortical and subcortical pathway maturation, such as the nigrostriatal pathway, may increase risk for neuropsychiatric disorders (e.g., mood disorders, cognitive dysfunction, brain connectivity dysfunction). However, additional studies are needed to confirm this relationship between late gestational hypoxia and human neuropsychiatric disorders. It is noteworthy that CIH exposure during the last five days of gestation had long-term adverse effects on the offspring, while it did not affect the placentation process which occurs at the beginning of pregnancy, and had minimal impact on gestational outcomes. These novel findings demonstrate that gestational stressors can result in long-term adverse effects in offspring even in the absence of detectable pregnancy or neonatal complications, especially if they occur during critical embryological developmental periods.

Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s13293-022-00463-x>.

Additional file 1: Table S1. Summary of offspring total and categorized ultrasonic vocalizations. **Table S2.** Two-Way ANOVA results for call types and latency to first call.

Acknowledgements

We would like to acknowledge UNTHSC Office of Vice President of Research for funding these studies.

Author contributions

ENW conducted experiments, analyzed data, and wrote the manuscript. SM conducted experiments and analyzed data. JLB analyzed and interpreted data and contributed to writing the manuscript. JG, NR, RE, OF, OO, and SC conducted experiments and assisted with tissue collection and processing. SG and RLC designed the study, analyzed data, interpreted the findings, and edited the manuscript. RLC, JLB, ENW, and SG were major contributors in writing the manuscript. All authors read and approved the final manuscript.

Funding

This study was supported by NIH R01 NS0091359 and UNTHSC Seed grant funding to RLC, NIH R01 HL146562-02S1 to SG, AHA 22PRE-900431 to JG, AHA 22POST-903250 to JB, and NIH T32 AG020494 to SM.

Availability of data and materials

The datasets used and/or analyzed during the current study are available from the corresponding author upon reasonable request.

Declarations

Ethics approval and consent to participate

This study was performed under IACUC protocol approved by the University of North Texas Health Science Center.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Author details

¹Department of Physiology and Anatomy, University of North Texas Health Science Center, Fort Worth, TX 76107, USA. ²Texas College of Osteopathic Medicine, University of North Texas Health Science Center, Fort Worth, TX 76107, USA. ³Department of Pharmaceutical Sciences, UNT System College of Pharmacy, School of Pharmacy, University of North Texas Health Science Center, 3500 Camp Bowie Boulevard, Fort Worth, TX 76107, USA. ⁴Present Address: Department of Basic Sciences, Lawrence D. Longo, MD Center for Perinatal Biology, Loma Linda University School of Medicine, Loma Linda, CA 92350, USA.

Received: 20 July 2022 Accepted: 22 September 2022

Published online: 30 September 2022

References

- Tong W, Giussani DA. Preeclampsia link to gestational hypoxia. *J Dev Orig Health Dis.* 2019;10(3):322–33. <https://doi.org/10.1017/S204017441900014X>.
- Almendros I, Martinez-Ros P, Farre N, Rubio-Zaragoza M, Torres M, Gutierrez-Bautista AJ, et al. Placental oxygen transfer reduces hypoxia-reoxygenation swings in fetal blood in a sheep model of gestational sleep apnea. *J Appl Physiol* (1985). 2019;127(3):745–52. <https://doi.org/10.1152/jappphysiol.00303.2019>.
- Jang EA, Longo LD, Goyal R. Antenatal maternal hypoxia: criterion for fetal growth restriction in rodents. *Front Physiol.* 2015;6:176. <https://doi.org/10.3389/fphys.2015.00176>.
- Ravishankar S, Bourjeily G, Lambert-Messerlian G, He M, De Paep ME, Gundogan F. Evidence of placental hypoxia in maternal sleep-disordered breathing. *Pediatr Dev Pathol.* 2015;18(5):380–6. <https://doi.org/10.2350/15-06-1647-OA.1>.
- Nalivaeva NN, Turner AJ, Zhuravin IA. Role of prenatal hypoxia in brain development, cognitive functions, and neurodegeneration. *Front Neurosci.* 2018;12:825. <https://doi.org/10.3389/fnins.2018.00825>.
- Facco FL, Parker CB, Reddy UM, Silver RM, Koch MA, Louis JM, et al. Association between sleep-disordered breathing and hypertensive disorders of pregnancy and gestational diabetes mellitus. *Obstet Gynecol.* 2017;129(1):31–41. <https://doi.org/10.1097/aog.0000000000001805>.
- Pien GW, Pack AI, Jackson N, Maislin G, Macones GA, Schwab RJ. Risk factors for sleep-disordered breathing in pregnancy. *Thorax.* 2014;69(4):371–7. <https://doi.org/10.1136/thoraxjnl-2012-202718>.
- Azevedo PN, Zanirati G, Venturin GT, Schu GG, Duran-Carabali LE, Odorczyk FK, et al. Long-term changes in metabolic brain network drive memory impairments in rats following neonatal hypoxia-ischemia. *Neurobiol Learn Mem.* 2020;171:1095–9564 (Electronic):107207; <https://doi.org/10.1016/j.nlm.2020.107207>.
- Zhuravin IA, Dubrovskaya NM, Vasilev DS, Postnikova TY, Zaitsev AV. Prenatal hypoxia produces memory deficits associated with impairment of long-term synaptic plasticity in young rats. *Neurobiol Learn Mem.* 2019;164:1095–9564 (Electronic):107066; doi:<https://doi.org/10.1016/j.nlm.2019.107066>.
- Yang SN, Huang CB, Yang CH, Lai MC, Chen WF, Wang CL, et al. Impaired SynGAP expression and long-term spatial learning and memory in

- hippocampal CA1 area from rats previously exposed to perinatal hypoxia-induced insults: beneficial effects of A68930. *Neurosci Lett*. 2004;371(1):73–8. <https://doi.org/10.1016/j.neulet.2004.08.044>.
11. Sailaja K, Gopinath G. Developing substantia nigra in human: a qualitative study. *Dev Neurosci*. 1994;16(1–2):44–52. <https://doi.org/10.1159/000112087>.
 12. Aubert I, Brana C, Pellevoisin C, Giros B, Caille I, Carles D, et al. Molecular anatomy of the development of the human substantia nigra. *J Comp Neurol*. 1997;379(1):72–87. [https://doi.org/10.1002/\(sici\)1096-9861\(19970303\)379:1<72::aid-cne5>3.0.co;2-f](https://doi.org/10.1002/(sici)1096-9861(19970303)379:1<72::aid-cne5>3.0.co;2-f).
 13. Almqvist PM, Åkesson E, Wahlberg LU, Pschera H, Seiger Å, Sundström E. First trimester development of the human nigrostriatal dopamine system. *Exp Neurol*. 1996;139(2):227–37. <https://doi.org/10.1006/exnr.1996.0096>.
 14. Chevassus-au-Louis N, Baraban SC, Gaiarsa JL, Ben-Ari Y. Cortical malformations and epilepsy: new insights from animal models. *Epilepsia*. 1999;40(7):811–21. <https://doi.org/10.1111/j.1528-1157.1999.tb00786.x>.
 15. Kim EH, Yum MS, Lee M, Kim EJ, Shim WH, Ko TS. A new rat model of epileptic spasms based on methylazoxymethanol-induced malformations of cortical development. *Front Neurol*. 2017;8:271. <https://doi.org/10.3389/fneur.2017.00271>.
 16. Rice D, Barone S Jr. Critical periods of vulnerability for the developing nervous system: evidence from humans and animal models. *Environ Health Perspect*. 2000;108(Suppl 3):511–33. <https://doi.org/10.1289/ehp.00108s3511>.
 17. Jung AB, Bennett JP Jr. Development of striatal dopaminergic function. I. Pre- and postnatal development of mRNAs and binding sites for striatal D1 (D1a) and D2 (D2a) receptors. *Brain Res Dev Brain Res*. 1996;94(2):109–20. [https://doi.org/10.1016/0165-3806\(96\)00033-8](https://doi.org/10.1016/0165-3806(96)00033-8).
 18. Kortheuer KH. A study of development stages of the corpus striatum of the human brain. Los Angeles, California: University of Southern California; 1929.
 19. Rickmann M, Wolff JR. Prenatal gliogenesis in the neopallium of the rat. *Adv Anat Embryol Cell Biol*. 1985;93(1):104. <https://doi.org/10.1007/978-3-642-70081-1>.
 20. Evans NP, Bellingham M, Robinson JE. Prenatal programming of neuroendocrine reproductive function. *Theriogenology*. 2016;86(1):340–8. <https://doi.org/10.1016/j.theriogenology.2016.04.047>.
 21. Hodes GE, Epperson CN. Sex differences in vulnerability and resilience to stress across the life span. *Biol Psychiatry*. 2019;86(6):421–32. <https://doi.org/10.1016/j.biopsych.2019.04.028>.
 22. Perez-Cereales S, Ramos-Ibeas P, Rizo D, Lonergan P, Bermejo-Alvarez P, Gutierrez-Adan A. Early sex-dependent differences in response to environmental stress. *Reproduction*. 2018;155(1):R39–51. <https://doi.org/10.1530/REP-17-0466>.
 23. Howerton CL, Bale TL. Targeted placental deletion of OGT recapitulates the prenatal stress phenotype including hypothalamic mitochondrial dysfunction. *Proc Natl Acad Sci U S A*. 2014;111(26):9639–44. <https://doi.org/10.1073/pnas.1401203111>.
 24. Mueller BR, Bale TL. Sex-specific programming of offspring emotionality after stress early in pregnancy. *J Neurosci*. 2008;28(36):9055–65. <https://doi.org/10.1523/JNEUROSCI.1424-08.2008>.
 25. Carpentier PA, Haditsch U, Braun AE, Cantu AV, Moon HM, Price RO, et al. Stereotypical alterations in cortical patterning are associated with maternal illness-induced placental dysfunction. *J Neurosci*. 2013;33(43):16874–88. <https://doi.org/10.1523/JNEUROSCI.4654-12.2013>.
 26. Braun AE, Carpentier PA, Babineau BA, Narayan AR, Kielhold ML, Moon HM, et al. "Females are not just 'protected' males": sex-specific vulnerabilities in placenta and brain after prenatal immune disruption. *eNeuro*. 2019;6:6. <https://doi.org/10.1523/ENEURO.0358-19.2019>.
 27. Eriksson JG, Kajantie E, Osmond C, Thornburg K, Barker DJ. Boys live dangerously in the womb. *Am J Hum Biol*. 2010;22(3):330–5. <https://doi.org/10.1002/ajhb.20995>.
 28. Sandman CA, Glynn LM, Davis EP. Is there a viability-vulnerability tradeoff? Sex differences in fetal programming. *J Psychosom Res*. 2013;75(4):327–35. <https://doi.org/10.1016/j.jpsychores.2013.07.009>.
 29. Behlen JC, Lau CH, Li Y, Dhagat P, Stanley JA, Rodrigues Hoffman A, et al. Gestational exposure to ultrafine particles reveals sex- and dose-specific changes in offspring birth outcomes, placental morphology, and gene networks. *Toxicol Sci*. 2021;184(2):204–13. <https://doi.org/10.1093/toxsci/kfab118>.
 30. Gumusoglu SB, Chilukuri ASS, Hing BWQ, Scroggins SM, Kundu S, Sandgren JA, et al. Altered offspring neurodevelopment in an arginine vasopressin preeclampsia model. *Transl Psychiatry*. 2021;11(1):79. <https://doi.org/10.1038/s41398-021-01205-0>.
 31. Beery AK, Zucker I. Sex bias in neuroscience and biomedical research. *Neurosci Biobehav Rev*. 2011;35(3):565–72. <https://doi.org/10.1016/j.neubiorev.2010.07.002>.
 32. Shansky RM, Woolley CS. Considering sex as a biological variable will be valuable for neuroscience research. *J Neurosci*. 2016;36(47):11817–22. <https://doi.org/10.1523/JNEUROSCI.1390-16.2016>.
 33. Coiro P, Pollak DD. Sex and gender bias in the experimental neurosciences: the case of the maternal immune activation model. *Transl Psychiatry*. 2019;9:1–90. <https://doi.org/10.1038/s41398-019-0423-8>.
 34. Ciucci MR, Ahrens AM, Ma ST, Kane JR, Windham EB, Woodlee MT, et al. Reduction of dopamine synaptic activity: degradation of 50-kHz ultrasonic vocalization in rats. *Behav Neurosci*. 2009;123(2):328–36. <https://doi.org/10.1037/a0014593>.
 35. Ciucci MR, Ma ST, Fox C, Kane JR, Ramig LO, Schallert T. Qualitative changes in ultrasonic vocalization in rats after unilateral dopamine depletion or haloperidol: a preliminary study. *Behav Brain Res*. 2007;182(2):284–9. <https://doi.org/10.1016/j.bbr.2007.02.020>.
 36. Grant LM, Kelm-Nelson CA, Hilby BL, Blue KV, Paul Rajamanickam ES, Pultorak JD, et al. Evidence for early and progressive ultrasonic vocalization and oromotor deficits in a PINK1 gene knockout rat model of Parkinson's disease. *J Neurosci Res*. 2015;93(11):1713–27. <https://doi.org/10.1002/jnr.23625>.
 37. Olszynski KH, Polowy R, Wardak AD, Grymanowska AW, Filipkowski RK. Increased vocalization of rats in response to ultrasonic playback as a sign of hypervigilance following fear conditioning. *Brain Sci*. 2021;11:8. <https://doi.org/10.3390/brainsci11080970>.
 38. Panksepp J. Affective consciousness: core emotional feelings in animals and humans. *Conscious Cogn*. 2005;14(1):30–80. <https://doi.org/10.1016/j.concog.2004.10.004>.
 39. Berz AC, Wöhr M, Schwarting RKW. Response calls evoked by playback of natural 50-kHz ultrasonic vocalizations in rats. *Front Behav Neurosci*. 2021;15: 812142. <https://doi.org/10.3389/fnbeh.2021.812142>.
 40. Mahrt EJ, Perkel DJ, Tong L, Rubel EW, Portfors CV. Engineered deafness reveals that mouse courtship vocalizations do not require auditory experience. *J Neurosci*. 2013;33(13):5573–83. <https://doi.org/10.1523/jneurosci.5054-12.2013>.
 41. Wright JM, Gourdon JC, Clarke PB. Identification of multiple call categories within the rich repertoire of adult rat 50-kHz ultrasonic vocalizations: effects of amphetamine and social context. *Psychopharmacology*. 2010;211(1):1–13. <https://doi.org/10.1007/s00213-010-1859-y>.
 42. Sewell GDS. Ultrasound and mating behaviour in rodents with some observations on other behavioural situations. *J Zool*. 2009;168(2):149–64. <https://doi.org/10.1111/j.1469-7998.1972.tb01345.x>.
 43. Burgdorf J, Kroes RA, Moskal JR, Pfau JG, Brudzynski SM, Panksepp J. Ultrasonic vocalizations of rats (*Rattus norvegicus*) during mating, play, and aggression: behavioral concomitants, relationship to reward, and self-administration of playback. *J Comp Psychol*. 2008;122(4):357–67. <https://doi.org/10.1037/a0012889>.
 44. McGinnis MY, Vakulenko M. Characterization of 50-kHz ultrasonic vocalizations in male and female rats. *Physiol Behav*. 2003;80(1):81–8. [https://doi.org/10.1016/s0031-9384\(03\)00227-0](https://doi.org/10.1016/s0031-9384(03)00227-0).
 45. Chabout J, Serreau P, Ey E, Bellier L, Aubin T, Bourgeron T, et al. Adult male mice emit context-specific ultrasonic vocalizations that are modulated by prior isolation or group rearing environment. *PLoS ONE*. 2012;7(1): e29401. <https://doi.org/10.1371/journal.pone.0029401>.
 46. Portfors CV. Types and functions of ultrasonic vocalizations in laboratory rats and mice. *J Am Assoc Lab Anim Sci*. 2007;46(1):28–34.
 47. Wöhr M, Schwarting RK. Ultrasonic communication in rats: can playback of 50-kHz calls induce approach behavior? *PLoS ONE*. 2007;2(12): e1365. <https://doi.org/10.1371/journal.pone.0001365>.
 48. Knutson B, Burgdorf J, Panksepp J. Anticipation of play elicits high-frequency ultrasonic vocalizations in young rats. *J Comp Psychol*. 1998;112(1):65–73. <https://doi.org/10.1037/0735-7036.112.1.65>.

49. Wöhr M, Houx B, Schwarting RK, Spruijt B. Effects of experience and context on 50-kHz vocalizations in rats. *Physiol Behav.* 2008;93(4–5):766–76. <https://doi.org/10.1016/j.physbeh.2007.11.031>.
50. Wang H, Liang S, Burgdorf J, Wess J, Yeomans J. Ultrasonic vocalizations induced by sex and amphetamine in M2, M4, M5 muscarinic and D2 dopamine receptor knockout mice. *PLoS ONE.* 2008;3(4): e1893. <https://doi.org/10.1371/journal.pone.0001893>.
51. Hanson JL, Hurlley LM. Female presence and estrous state influence mouse ultrasonic courtship vocalizations. *PLoS ONE.* 2012;7(7): e40782. <https://doi.org/10.1371/journal.pone.0040782>.
52. Matsumoto YK, Okanoya K. Phase-specific vocalizations of male mice at the initial encounter during the courtship sequence. *PLoS ONE.* 2016;11(2): e0147102. <https://doi.org/10.1371/journal.pone.0147102>.
53. Schwarting RKW. Ultrasonic vocalization in juvenile and adult male rats: a comparison among stocks. *Physiol Behav.* 2018;191:1–11. <https://doi.org/10.1016/j.physbeh.2018.03.023>.
54. Brudzynski SM, Pniak A. Social contacts and production of 50-kHz short ultrasonic calls in adult rats. *J Comp Psychol.* 2002;116(1):73–82. <https://doi.org/10.1037/0735-7036.116.1.73>.
55. Newar SL, Bowman J. Think before they squeak: vocalizations of the squirrel family. *Front Ecol Evol.* 2020;8:193. <https://doi.org/10.3389/fevo.2020.00193>.
56. Blanchard RJ, Blanchard DC, Agullana R, Weiss SM. Twenty-two kHz alarm cries to presentation of a predator, by laboratory rats living in visible burrow systems. *Physiol Behav.* 1991;50(5):967–72. [https://doi.org/10.1016/0031-9384\(91\)90423-l](https://doi.org/10.1016/0031-9384(91)90423-l).
57. Fendt M, Brosch M, Wernecke KEA, Willadsen M, Wöhr M. Predator odour but not TMT induces 22-kHz ultrasonic vocalizations in rats that lead to defensive behaviours in conspecifics upon replay. *Sci Rep.* 2018;8(1):11041. <https://doi.org/10.1038/s41598-018-28927-4>.
58. Chan CS, Guzman JN, Ilijic E, Mercer JN, Rick C, Tkatch T, et al. "Rejuvenation" protects neurons in mouse models of Parkinson's disease. *Nature.* 2007;447(7148):1081–6.
59. Osikoya O, Jaini PA, Nguyen A, Valdes M, Gouloupoulos S. Effects of low-dose aspirin on maternal blood pressure and vascular function in an experimental model of gestational hypertension. *Pharmacol Res.* 2017;120:267–78. <https://doi.org/10.1016/j.phrs.2017.04.012>.
60. Snyder B, Shell B, Cunningham JT, Cunningham RL. Chronic intermittent hypoxia induces oxidative stress and inflammation in brain regions associated with early-stage neurodegeneration. *Physiol Rep.* 2017;5:9. <https://doi.org/10.14814/phy2.13258>.
61. Snyder B, Duong P, Tenkorang M, Wilson EN, Cunningham RL. Rat strain and housing conditions alter oxidative stress and hormone responses to chronic intermittent hypoxia. *Front Physiol.* 2018;9:1554. <https://doi.org/10.3389/fphys.2018.01554>.
62. Snyder B, Duong P, Trieu J, Cunningham RL. Androgens modulate chronic intermittent hypoxia effects on brain and behavior. *Horm Behav.* 2018;106:62–73. <https://doi.org/10.1016/j.yhbeh.2018.09.005>.
63. Wilson EN, Anderson M, Snyder B, Duong P, Trieu J, Schreihof DA, et al. Chronic intermittent hypoxia induces hormonal and male sexual behavioral changes: hypoxia as an advancer of aging. *Physiol Behav.* 2018;189:64–73. <https://doi.org/10.1016/j.physbeh.2018.03.007>.
64. Johnson AM, Doll EJ, Grant LM, Ringel L, Shier JN, Ciucci MR. Targeted training of ultrasonic vocalizations in aged and Parkinsonian rats. *J Vis Exp.* 2011;54. <https://doi.org/10.3791/2835>.
65. Neunuebel JP, Taylor AL, Arthur BJ, Egnor SE. Female mice ultrasonically interact with males during courtship displays. *Elife.* 2015;4. <https://doi.org/10.7554/eLife.06203>.
66. Matochik JA, White NR, Barfield RJ. Variations in scent marking and ultrasonic vocalizations by Long-Evans rats across the estrous cycle. *Physiol Behav.* 1992;51(4):783–6. [https://doi.org/10.1016/0031-9384\(92\)90116-j](https://doi.org/10.1016/0031-9384(92)90116-j).
67. Warren MR, Spurrier MS, Roth ED, Neunuebel JP. Sex differences in vocal communication of freely interacting adult mice depend upon behavioral context. *PLoS ONE.* 2018;13(9): e0204527. <https://doi.org/10.1371/journal.pone.0204527>.
68. Paxinos G, Watson C. *The rat brain in stereotaxic coordinates*. 4th ed. San Diego: Academic Press; 1998.
69. Garza-Contreras J, Duong P, Snyder BD, Schreihof DA, Cunningham RL. Presence of androgen receptor variant in neuronal lipid rafts. *eNeuro.* 2017;4:4. <https://doi.org/10.1523/ENEURO.0109-17.2017>.
70. Zhang Z, Larner SF, Liu MC, Zheng W, Hayes RL, Wang KK. Multiple alphaspectrin breakdown products distinguish calpain and caspase dominated necrotic and apoptotic cell death pathways. *Apoptosis.* 2009;14(11):1289–98. <https://doi.org/10.1007/s10495-009-0405-z>.
71. Czogalla A, Sikorski AF. Spectrin and calpain: a "target" and a "sniper" in the pathology of neuronal cells. *Cell Mol Life Sci.* 2005;62(17):1913–24. <https://doi.org/10.1007/s00018-005-5097-0>.
72. Kelm-Nelson CA, Brauer AFL, Barth KJ, Lake JM, Sinnen MLK, Stehula FJ, et al. Characterization of early-onset motor deficits in the Pink1-/- mouse model of Parkinson disease. *Brain Res.* 2018;1680:1–12. <https://doi.org/10.1016/j.brainres.2017.12.002>.
73. Grant LM, Barnett DG, Doll EJ, Levenson G, Ciucci M. Relationships among rat ultrasonic vocalizations, behavioral measures of striatal dopamine loss, and striatal tyrosine hydroxylase immunoreactivity at acute and chronic time points following unilateral 6-hydroxydopamine-induced dopamine depletion. *Behav Brain Res.* 2015;291:361–71. <https://doi.org/10.1016/j.bbr.2015.05.042>.
74. Clancy B, Finlay BL, Darlington RB, Anand KJ. Extrapolating brain development from experimental species to humans. *Neurotoxicology.* 2007;28(5):931–7. <https://doi.org/10.1016/j.neuro.2007.01.014>.
75. Sengupta P. The laboratory rat: relating its age with human's. *Int J Prev Med.* 2013;4(6):624–30.
76. Semple BD, Blomgren K, Gimlin K, Ferriero DM, Noble-Haesslein LJ. Brain development in rodents and humans: Identifying benchmarks of maturation and vulnerability to injury across species. *Prog Neurobiol.* 2013;106–107:1–16. <https://doi.org/10.1016/j.pneurobio.2013.04.001>.
77. Bayer SA, Altman J, Russo RJ, Zhang X. Timetables of neurogenesis in the human brain based on experimentally determined patterns in the rat. *Neurotoxicology.* 1993;14(1):83–144.
78. Ferriero DM. Neonatal brain injury. *N Engl J Med.* 2004;351(19):1985–95. <https://doi.org/10.1056/NEJMra041996>.
79. Mandic-Maravic V, Grujicic R, Milutinovic L, Munjiza-Jovanovic A, Pejovic-Milovancevic M. Dopamine in autism spectrum disorders-focus on D2/D3 partial agonists and their possible use in treatment. *Front Psychiatry.* 2021;12: 787097. <https://doi.org/10.3389/fpsy.2021.787097>.
80. Romanos M, Weise D, Schliesser M, Schecklmann N, Löffler J, Warnke A, et al. Structural abnormality of the substantia nigra in children with attention-deficit hyperactivity disorder. *J Psychiatry Neurosci.* 2010;35(1):55–8. <https://doi.org/10.1503/jpn.090044>.
81. Bouchet CA, Miner MA, Loetz EC, Rosberg AJ, Hake HS, Farmer CE, et al. Activation of nigrostriatal dopamine neurons during fear extinction prevents the renewal of fear. *Neuropsychopharmacology.* 2018;43(3):665–72. <https://doi.org/10.1038/npp.2017.235>.
82. Ugrumov M. Development of early diagnosis of Parkinson's disease: Illusion or reality? *CNS Neurosci Ther.* 2020;26(10):997–1009. <https://doi.org/10.1111/cns.13429>.
83. Gumusoglu SB, Chilukuri ASS, Santillan DA, Santillan MK, Stevens HE. Neurodevelopmental outcomes of prenatal preeclampsia exposure. *Trends Neurosci.* 2020;43(4):253–68. <https://doi.org/10.1016/j.tins.2020.02.003>.
84. Korzeniewski SJ, Slaughter J, Lenski M, Haak P, Paneth N. The complex aetiology of cerebral palsy. *Nat Rev Neurol.* 2018;14(9):528–43. <https://doi.org/10.1038/s41582-018-0043-6>.
85. Dachew BA, Mamun A, Maravilla JC, Alati R. Association between hypertensive disorders of pregnancy and the development of offspring mental and behavioural problems: a systematic review and meta-analysis. *Psychiatry Res.* 2018;260:458–67. <https://doi.org/10.1016/j.psychres.2017.12.027>.
86. Dachew BA, Mamun A, Maravilla JC, Alati R. Pre-eclampsia and the risk of autism-spectrum disorder in offspring: meta-analysis. *Br J Psychiatry.* 2018;212(3):142–7. <https://doi.org/10.1192/bjp.2017.27>.
87. Maher GM, O'Keefe GW, Kearney PM, Kenny LC, Dinan TG, Mattsson M, et al. Association of hypertensive disorders of pregnancy with risk of neurodevelopmental disorders in offspring: a systematic review and meta-analysis. *JAMA Psychiat.* 2018;75(8):809–19. <https://doi.org/10.1001/jamapsychiatry.2018.0854>.
88. Dachew BA, Scott JG, Mamun A, Alati R. Pre-eclampsia and the risk of attention-deficit/hyperactivity disorder in offspring: findings from the

- ALSPAC birth cohort study. *Psychiatry Res.* 2019;272:392–7. <https://doi.org/10.1016/j.psychres.2018.12.123>.
89. Walker CK, Krakowiak P, Baker A, Hansen RL, Ozonoff S, Hertz-Picciotto I. Preeclampsia, placental insufficiency, and autism spectrum disorder or developmental delay. *JAMA Pediatr.* 2015;169(2):154–62. <https://doi.org/10.1001/jamapediatrics.2014.2645>.
 90. Dachew BA, Scott JG, Mamun A, Alati R. Hypertensive disorders of pregnancy and the risk of anxiety disorders in adolescence: findings from the Avon longitudinal study of parents and children. *J Psychiatr Res.* 2019;110:159–65. <https://doi.org/10.1016/j.jpsychires.2019.01.001>.
 91. Nahum Sacks K, Friger M, Shoham-Vardi I, Sergienko R, Spiegel E, Landau D, et al. Long-term neuropsychiatric morbidity in children exposed prenatally to preeclampsia. *Early Hum Dev.* 2019;130:96–100. <https://doi.org/10.1016/j.earlhumdev.2019.01.016>.
 92. Lahti-Pulkkinen M, Girchenko P, Tuovinen S, Sammallahti S, Reynolds RM, Lahti J, et al. Maternal hypertensive pregnancy disorders and mental disorders in children. *Hypertension.* 2020;75(6):1429–38. <https://doi.org/10.1161/HYPERTENSIONAHA.119.14140>.
 93. Sun BZ, Moster D, Harmon QE, Wilcox AJ. Association of preeclampsia in term births with neurodevelopmental disorders in offspring. *JAMA Psychiatr.* 2020;77(8):823–9. <https://doi.org/10.1001/jamapsychiatry.2020.0306>.
 94. Davis EF, Lazdam M, Lewandowski AJ, Worton SA, Kelly B, Kenworthy Y, et al. Cardiovascular risk factors in children and young adults born to preeclamptic pregnancies: a systematic review. *Pediatrics.* 2012;129(6):e1552–61. <https://doi.org/10.1542/peds.2011-3093>.
 95. Tauman R, Zuk L, Uliei-Sibony S, Ascher-Landsberg J, Katsav S, Farber M, et al. The effect of maternal sleep-disordered breathing on the infant's neurodevelopment. *Am J Obstet Gynecol.* 2015;212(5):656 e1–657. <https://doi.org/10.1016/j.ajog.2015.01.001>.
 96. Bin YS, Cistulli PA, Roberts CL, Ford JB. Childhood health and educational outcomes associated with maternal sleep apnea: a population record-linkage study. *Sleep.* 2017;40:11. <https://doi.org/10.1093/sleep/zsx158>.
 97. Kanno K, Kikusui T. Effect of sociosexual experience and aging on number of courtship ultrasonic vocalizations in male mice. *Zoolog Sci.* 2018;35(3):208–14. <https://doi.org/10.2108/zs170175>.
 98. Heckman JJ, Provill R, Heckman GJ, Azarfar A, Celikel T, Englitz B. High-precision spatial localization of mouse vocalizations during social interaction. *Sci Rep.* 2017;7(1):3017. <https://doi.org/10.1038/s41598-017-02954-z>.
 99. Zala SM, Reitschmidt D, Noll A, Balazs P, Penn DJ. Sex-dependent modulation of ultrasonic vocalizations in house mice (*Mus musculus musculus*). *PLoS ONE.* 2017;12(12): e0188647. <https://doi.org/10.1371/journal.pone.0188647>.
 100. Hammerschmidt K, Radyushkin K, Ehrenreich H, Fischer J. The structure and usage of female and male mouse ultrasonic vocalizations reveal only minor differences. *PLoS ONE.* 2012;7(7): e41133. <https://doi.org/10.1371/journal.pone.0041133>.
 101. D'Amato FR, Moles A. Ultrasonic vocalizations as an index of social memory in female mice. *Behav Neurosci.* 2001;115(4):834–40. <https://doi.org/10.1037//0735-7044.115.4.834>.
 102. Bishnoi IR, Ossenkopp KP, Kavaliers M. Sex and age differences in locomotor and anxiety-like behaviors in rats: from adolescence to adulthood. *Dev Psychobiol.* 2021;63(3):496–511. <https://doi.org/10.1002/dev.22037>.
 103. Archer J. Rodent sex differences in emotional and related behavior. *Behav Biol.* 1975;14(4):451–79. [https://doi.org/10.1016/s0091-6773\(75\)90636-7](https://doi.org/10.1016/s0091-6773(75)90636-7).
 104. Beatty WW, Fessler RG. Ontogeny of sex differences in open-field behavior and sensitivity to electric shock in the rat. *Physiol Behav.* 1976;16(4):413–7. [https://doi.org/10.1016/0031-9384\(76\)90319-x](https://doi.org/10.1016/0031-9384(76)90319-x).
 105. Masur J, Schutz MT, Boerngen R. Gender differences in open-field behavior as a function of age. *Dev Psychobiol.* 1980;13(2):107–10. <https://doi.org/10.1002/dev.420130202>.
 106. Acevedo MB, Pautassi RM, Spear NE, Spear LP. Age-dependent effects of stress on ethanol-induced motor activity in rats. *Psychopharmacology.* 2013;230(3):389–98. <https://doi.org/10.1007/s00213-013-3163-0>.
 107. Candland DK, Campbell BA. Development of fear in the rat as measured by behavior in the open field. *J Comp Physiol Psychol.* 1962;55(4):593–6. <https://doi.org/10.1037/h0047206>.
 108. Marin MT, Zancheta R, Paro AH, Possi AP, Cruz FC, Planeta CS. Comparison of caffeine-induced locomotor activity between adolescent and adult rats. *Eur J Pharmacol.* 2011;660(2–3):363–7. <https://doi.org/10.1016/j.ejphar.2011.03.052>.
 109. Lynn DA, Brown GR. The ontogeny of exploratory behavior in male and female adolescent rats (*Rattus norvegicus*). *Dev Psychobiol.* 2009;51(6):513–20. <https://doi.org/10.1002/dev.20386>.
 110. McDougall SA, Park GI, Ramirez GI, Gomez V, Adame BC, Crawford CA. Sex-dependent changes in ketamine-induced locomotor activity and ketamine pharmacokinetics in preweaning, adolescent, and adult rats. *Eur Neuropsychopharmacol.* 2019;29(6):740–55. <https://doi.org/10.1016/j.euroneuro.2019.03.013>.
 111. Crawford CA, Moran AE, Baum TJ, Apodaca MG, Montejano NR, Park GI, et al. Effects of monoamine depletion on the ketamine-induced locomotor activity of preweaning, adolescent, and adult rats: Sex and age differences. *Behav Brain Res.* 2020;379: 112267. <https://doi.org/10.1016/j.bbr.2019.112267>.
 112. Knight P, Chellian R, Wilson R, Behnood-Rod A, Panunzio S, Bruijnzeel AW. Sex differences in the elevated plus-maze test and large open field test in adult Wistar rats. *Pharmacol Biochem Behav.* 2021;204: 173168. <https://doi.org/10.1016/j.pbb.2021.173168>.
 113. Gogokhia N, Japaridze N, Tizabi Y, Pataraya L, Zhvania MG. Gender differences in anxiety response to high intensity white noise in rats. *Neurosci Lett.* 2021;742: 135543. <https://doi.org/10.1016/j.neulet.2020.135543>.
 114. Scholl JL, Afzal A, Fox LC, Watt MJ, Forster GL. Sex differences in anxiety-like behaviors in rats. *Physiol Behav.* 2019;211: 112670. <https://doi.org/10.1016/j.physbeh.2019.112670>.
 115. Gozal D, Reeves SR, Row BW, Neville JJ, Guo SZ, Lipton AJ. Respiratory effects of gestational intermittent hypoxia in the developing rat. *Am J Respir Crit Care Med.* 2003;167(11):1540–7. <https://doi.org/10.1164/rccm.200208-963OC>.
 116. Chen L, Zadi ZH, Zhang J, Scharf SM, Pae EK. Intermittent hypoxia in utero damages postnatal growth and cardiovascular function in rats. *J Appl Physiol* (1985). 2018;124(4):821–30. <https://doi.org/10.1152/japplphysiol.01066.2016>.
 117. Vanderplow AM, Kermath BA, Bernhardt CR, Gums KT, Seablom EN, Radcliff AB, et al. A feature of maternal sleep apnea during gestation causes autism-relevant neuronal and behavioral phenotypes in offspring. *PLoS Biol.* 2022;20(2): e3001502. <https://doi.org/10.1371/journal.pbio.3001502>.
 118. Su YM, Lv GR, Xie JX, Wang ZH, Lin HT. Maternal hypoxia increases the susceptibility of adult rat male offspring to high-fat diet-induced nonalcoholic fatty liver disease. *Endocrinology.* 2013;154(11):4377–87. <https://doi.org/10.1210/en.2012-1683>.
 119. Wang Z, Huang Z, Lu G, Lin L, Ferrari M. Hypoxia during pregnancy in rats leads to early morphological changes of atherosclerosis in adult offspring. *Am J Physiol Heart Circ Physiol.* 2009;296(5):H1321–8. <https://doi.org/10.1152/ajpheart.00440.2008>.
 120. Nuzzo AM, Camm EJ, Sferruzzi-Perrin AN, Ashmore TJ, Yung HW, Cindrova-Davies T, et al. Placental adaptation to early-onset hypoxic pregnancy and mitochondria-targeted antioxidant therapy in a rodent model. *Am J Pathol.* 2018;188(12):2704–16. <https://doi.org/10.1016/j.ajpath.2018.07.027>.
 121. Hellgren KT, Premanandhan H, Quinn CJ, Trafford AW, Galli GLJ. Sex-dependent effects of developmental hypoxia on cardiac mitochondria from adult murine offspring. *Free Radic Biol Med.* 2021;162:490–9. <https://doi.org/10.1016/j.freeradbiomed.2020.11.004>.
 122. Hula N, Spaans F, Vu J, Quon A, Kirschenman R, Cooke CM, et al. Placental treatment improves cardiac tolerance to ischemia/reperfusion insult in adult male and female offspring exposed to prenatal hypoxia. *Pharmacol Res.* 2021;165: 105461. <https://doi.org/10.1016/j.phrs.2021.105461>.
 123. Rueda-Clausen CF, Stanley JL, Thambiraj DF, Poudel R, Davidge ST, Baker PN. Effect of prenatal hypoxia in transgenic mouse models of preeclampsia and fetal growth restriction. *Reprod Sci.* 2014;21(4):492–502. <https://doi.org/10.1177/1933719113503401>.
 124. Cahill LS, Hoggarth J, Lerch JP, Seed M, Macgowan CK, Sled JG. Fetal brain sparing in a mouse model of chronic maternal hypoxia. *J Cereb Blood Flow Metab.* 2019;39(6):1172–84. <https://doi.org/10.1177/0271678X17750324>.

125. Golan H, Kashtutsky I, Hallak M, Sorokin Y, Huleihel M. Maternal hypoxia during pregnancy delays the development of motor reflexes in newborn mice. *Dev Neurosci*. 2004;26(1):24–9. <https://doi.org/10.1159/000080708>.
126. Crisancho AG, Gadra EC, Samba IM, Zhao C, Ouyang M, Magnitsky S, et al. Deficits in seizure threshold and other behaviors in adult mice without gross neuroanatomic injury after late gestation transient prenatal hypoxia. *Dev Neurosci*. 2022. <https://doi.org/10.1159/000524045>.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

